










Madera County – Chlamydia, Gonorrhea, and P&S Syphilis

Rates by Age Group (2002), Race/Ethnicity (2002), and Year



■ Female □ Male

— Female — Male

* Race data may be missing for a substantial number of cases. See the "Data Limitations" page for further information.

Note: Rates are per 100,000 population.

Source: California Department of Health Services, STD Control Branch



Madera County – Chlamydia, Gonorrhea, and Early Syphilis

Cases and Rates Tables for 2002

| Gender & Age Group | Chlamydia | | | Gonorrhea | | | P&S Syphilis | | | Early Latent Syphilis | | | Population ² | |
|---------------------|-----------|----------------------|---------|-----------|----------------------|-------|--------------|----------------------|------|-----------------------|----------------------|------|-------------------------|----------------------|
| | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Number | Percent ¹ |
| COUNTY TOTAL | 423 | 100.0% | 320.9 | 54 | 100.0% | 41.0 | 1 | 100.0% | 0.8 | 0 | 0.0% | 0.0 | 135,695 | 100.0% |
| Female Total | 350 | 82.7% | 503.5 | 38 | 70.4% | 54.7 | 1 | 100.0% | 1.4 | 0 | 0.0% | 0.0 | 69,518 | 51.2% |
| Ages | | | | | | | | | | | | | | |
| 0 - 9 | 1 | 0.3% | 9.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 11,159 | 16.1% |
| 10 - 14 | 7 | 2.0% | 137.3 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 5,100 | 7.3% |
| 15 - 19 | 125 | 35.7% | 2,623.8 | 8 | 21.1% | 167.9 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 4,764 | 6.9% |
| 20 - 24 | 110 | 31.4% | 1,786.9 | 14 | 36.8% | 227.4 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 6,156 | 8.9% |
| 25 - 29 | 53 | 15.1% | 838.1 | 11 | 28.9% | 173.9 | 1 | 100.0% | 15.8 | 0 | 0.0% | 0.0 | 6,324 | 9.1% |
| 30 - 34 | 26 | 7.4% | 522.1 | 1 | 2.6% | 20.1 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 4,980 | 7.2% |
| 35 - 44 | 21 | 6.0% | 246.7 | 3 | 7.9% | 35.2 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 8,512 | 12.2% |
| 45+ | 7 | 2.0% | 31.1 | 1 | 2.6% | 4.4 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 22,523 | 32.4% |
| Not Specified | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | - | - |
| Male Total | 73 | 17.3% | 110.3 | 16 | 29.6% | 24.2 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 66,177 | 48.8% |
| Ages | | | | | | | | | | | | | | |
| 0 - 9 | 1 | 1.4% | 8.5 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 11,811 | 17.8% |
| 10 - 14 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 5,501 | 8.3% |
| 15 - 19 | 13 | 17.8% | 253.3 | 2 | 12.5% | 39.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 5,132 | 7.8% |
| 20 - 24 | 25 | 34.2% | 472.1 | 9 | 56.3% | 170.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 5,295 | 8.0% |
| 25 - 29 | 17 | 23.3% | 346.4 | 1 | 6.3% | 20.4 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 4,907 | 7.4% |
| 30 - 34 | 7 | 9.6% | 157.8 | 2 | 12.5% | 45.1 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 4,435 | 6.7% |
| 35 - 44 | 7 | 9.6% | 85.8 | 1 | 6.3% | 12.3 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 8,154 | 12.3% |
| 45+ | 3 | 4.1% | 14.3 | 1 | 6.3% | 4.8 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 20,942 | 31.6% |
| Not Specified | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | - | - |

| Gender & Race/Ethnicity | Chlamydia | | | Gonorrhea | | | P&S Syphilis | | | Early Latent Syphilis | | | Population ² | |
|-------------------------------|-----------|----------------------|---------|-----------|----------------------|------|--------------|----------------------|------|-----------------------|----------------------|------|-------------------------|----------------------|
| | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Cases | Percent ¹ | Rate | Number | Percent ¹ |
| COUNTY TOTAL | 423 | 100.0% | 320.9 | 54 | 100.0% | 41.0 | 1 | 100.0% | 0.8 | 0 | 0.0% | 0.0 | 135,695 | 100.0% |
| Female Total | 350 | 82.7% | 503.5 | 38 | 70.4% | 54.7 | 1 | 100.0% | 1.4 | 0 | 0.0% | 0.0 | 69,518 | 51.2% |
| American Indian/Alaska Native | 2 | 0.7% | 274.7 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 728 | 1.0% |
| Asian/Pacific Islander | 1 | 0.3% | 103.6 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 965 | 1.4% |
| Black/African-American | 14 | 4.7% | 331.8 | 1 | 3.7% | 23.7 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 4,220 | 6.1% |
| Latino/Hispanic | 259 | 87.5% | 1,011.2 | 23 | 85.2% | 89.8 | 1 | 100.0% | 3.9 | 0 | 0.0% | 0.0 | 25,614 | 36.8% |
| White | 20 | 6.8% | 52.6 | 3 | 11.1% | 7.9 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 37,991 | 54.6% |
| Not Specified | 54 | 15.4% | - | 11 | 28.9% | - | 0 | 0.0% | - | 0 | 0.0% | - | - | - |
| Male Total | 73 | 17.3% | 110.3 | 16 | 29.6% | 24.2 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 66,177 | 48.8% |
| American Indian/Alaska Native | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 612 | 0.9% |
| Asian/Pacific Islander | 1 | 1.4% | 99.2 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 1,008 | 1.5% |
| Black/African-American | 6 | 8.7% | 266.9 | 2 | 14.3% | 89.0 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 2,248 | 3.4% |
| Latino/Hispanic | 52 | 75.4% | 194.8 | 10 | 71.4% | 37.5 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 26,693 | 40.3% |
| White | 10 | 14.5% | 28.1 | 2 | 14.3% | 5.6 | 0 | 0.0% | 0.0 | 0 | 0.0% | 0.0 | 35,616 | 53.8% |
| Not Specified | 4 | 5.5% | - | 2 | 12.5% | - | 0 | 0.0% | - | 0 | 0.0% | - | - | - |

¹ Gender specific age groups and race/ethnicity percent calculations exclude "Not Specified" from the denominator.

² Overall county rates were calculated using the July 1 county population estimate from the Department of Finance.

(For Madera County, the July 1, 2002 estimate is 131,800).

Note: Rates are per 100,000 population.

| P&S Syphilis Males Cases | 2000 | | 2001 | | 2002 | |
|-----------------------------|------|------|------|------|------|------|
| | # | % | # | % | # | % |
| Total Males Cases | 0 | | 0 | | 0 | |
| ► MSM Cases | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |
| ►► HIV Positives among MSM | 0 | 0.0% | 0 | 0.0% | 0 | 0.0% |